

NEUTRINO MASS MODELS AND THE IMPLICATIONS OF A NON-ZERO REACTOR ANGLE

S.F.KING

*School of Physics and Astronomy,
University of Southampton,
Southampton SO17 1BJ, UK
E-mail: king@soton.ac.uk*

ABSTRACT

In this talk we survey some of the recent promising developments in the search for the theory behind neutrino mass and mixing, and indeed all fermion masses and mixing. The talk is organized in terms of a neutrino mass models decision tree according to which the answers to experimental questions provide sign posts to guide us through the maze of theoretical models eventually towards a complete theory of flavour and unification. We also discuss the theoretical implications of the measurement of a non-zero reactor angle, as hinted at by recent experimental measurements.

1. Introduction

It has been one of the long standing goals of theories of particle physics beyond the Standard Model (SM) to predict quark and lepton masses and mixings. With the discovery of neutrino mass and mixing, this quest has received a massive impetus. Indeed, perhaps the greatest advance in particle physics over the past decade has been the discovery of neutrino mass and mixing involving two large mixing angles commonly known as the atmospheric angle θ_{23} and the solar angle θ_{12} , while the remaining mixing angle θ_{13} , although unmeasured, is constrained to be relatively small. The largeness of the two large lepton mixing angles contrasts sharply with the smallness of the quark mixing angles, and this observation, together with the smallness of neutrino masses, provides new and tantalizing clues in the search for the origin of quark and lepton flavour. However, before trying to address such questions, it is worth recalling why neutrino mass forces us to go beyond the SM.

2. Why go beyond the Standard Model?

Neutrino mass is zero in the SM for three independent reasons:

1. There are no right-handed neutrinos ν_R .
2. There are only Higgs doublets of $SU(2)_L$.
3. There are only renormalizable terms.

In the SM these conditions all apply and so neutrinos are massless with ν_e, ν_μ, ν_τ distinguished by separate lepton numbers L_e, L_μ, L_τ . Neutrinos and antineutrinos are distinguished by total conserved lepton number $L = L_e + L_\mu + L_\tau$. To generate neutrino mass we must relax one or more of these conditions. For example, by adding right-handed neutrinos the Higgs mechanism of the Standard Model can give neutrinos the same type of mass as the electron mass or other charged lepton and quark masses. It is clear that the *status quo* of staying within the SM, as it is usually defined, is not an option, but in what direction should we go?

3. A decision tree

This talk will be organized according to the decision tree in Fig.1. Such a decision tree is clearly not unique (everyone can come up with her or his personal decision tree). The decision tree in Fig.1 contains key experimental questions (in blue) which serve as signposts along the way, leading in particular theoretical directions, starting from the top left hand corner with the question “LSND True or False?”




Figure 1: Neutrino mass models decision tree.

4. LSND True or False?

The results from MiniBOONE do not support the LSND result, but are consistent with the three active neutrino oscillation paradigm. If LSND were correct then this could imply either sterile neutrinos and/or CPT violation, or something more exotic.

For the remainder of this talk we shall assume that LSND is false, and focus on models without sterile neutrinos.

5. Dirac or Majorana?

Majorana neutrino masses are of the form $m_{LL}^\nu \bar{\nu}_L \nu_L^c$ where ν_L is a left-handed neutrino field and ν_L^c is the CP conjugate of a left-handed neutrino field, in other words a right-handed antineutrino field. Such Majorana masses are possible since both the neutrino and the antineutrino are electrically neutral. Such Majorana neutrino masses violate total lepton number L conservation, so the neutrino is equal to its own antiparticle. If we introduce right-handed neutrino fields then there are two sorts of additional neutrino mass terms that are possible. There are additional Majorana masses of the form $M_{RR}^\nu \bar{\nu}_R \nu_R^c$. In addition there are Dirac masses of the form $m_{LR}^\nu \bar{\nu}_L \nu_R$. Such Dirac mass terms conserve total lepton number L , but violate separate lepton numbers L_e, L_μ, L_τ . The question of “Dirac or Majorana?” is a key experimental question which could be decided by the experiments which measure neutrino masses directly.

6. What if Neutrinos are Dirac?

Introducing right-handed neutrinos ν_R into the SM (with zero Majorana mass) we can generate a Dirac neutrino mass from a coupling to the Higgs: $\lambda_\nu < H > \bar{\nu}_L \nu_R \equiv m_{LR}^\nu \bar{\nu}_L \nu_R$, where $< H > \approx 175$ GeV is the Higgs vacuum expectation value (VEV). A physical neutrino mass of $m_{LR}^\nu \approx 0.2$ eV implies $\lambda_\nu \approx 10^{-12}$. The question is why are such neutrino Yukawa couplings so small, even compared to the charged fermion Yukawa couplings? One possibility for small Dirac masses comes from the idea of extra dimensions motivated by theoretical attempts to extend the Standard Model to include gravity .

For the case of “flat” extra dimensions, “compactified” on circles of small radius R so that they are not normally observable, it has been suggested that right-handed neutrinos (but not the rest of the Standard Model particles) experience one or more of these extra dimensions ¹⁾. For example, for one extra dimension the right-handed neutrino wavefunction spreads out over the extra dimension R , leading to a suppressed Higgs interaction with the left-handed neutrino. The Dirac neutrino mass is therefore suppressed relative to the electron mass, and may be estimated as: $m_{LR}^\nu \sim \frac{M_{string}}{M_{Planck}} m_e$ where $M_{Planck} \sim 10^{19}$ GeV/ c^2 , and M_{string} is the string scale. Clearly low string scales, below the Planck scale, can lead to suppressed Dirac neutrino masses. Similar suppressions can be achieved with anisotropic compactifications ²⁾.

For the case of “warped” extra dimensions things are more complicated/interesting ³⁾. Typically there are two branes, a “Planck brane” and a “TeV brane”, with all the fermions and the Higgs in the “bulk” and having different “wavefunctions” which are

more or less strongly peaked on the TeV brane. The strength of the Yukawa coupling to the Higgs is determined by the overlap of a particular fermion wavefunction with the Higgs wavefunction, leading to exponentially suppressed Dirac masses. For example the Higgs and top quark wavefunctions are both strongly peaked on the TeV brane, leading to a large top quark mass, while the neutrino wavefunctions will be strongly peaked on the Planck brane leading to exponentially suppressed Dirac masses.

7. What if Neutrinos are Majorana?

We have already remarked that neutrinos, being electrically neutral, allow the possibility of Majorana neutrino masses. However such masses are forbidden in the SM since neutrinos form part of a lepton doublet L , and the Higgs field also forms a doublet H , and $SU(2)_L \times U(1)_Y$ gauge invariance forbids a Yukawa interaction like HLL . So, if we want to obtain Majorana masses, we must go beyond the SM.

One possibility is to introduce Higgs triplets Δ such that a Yukawa interaction like ΔLL is allowed. However the limit from the SM ρ parameter implies that the Higgs triplet should have a VEV $\langle \Delta \rangle < 8$ GeV. One big advantage is that the Higgs triplets may be discovered at the LHC and so this mechanism of neutrino mass generation is directly testable ⁴⁾.

Another possibility, originally suggested by Weinberg, is that neutrino Majorana masses originate from operators $HHLL$ involving two Higgs doublets and two lepton doublets, which, being higher order, must be suppressed by some large mass scale(s) M . When the Higgs doublets get their VEVs Majorana neutrino masses result: $m_{LL}^\nu = \lambda_\nu \langle H \rangle^2 / M$. This is nice because the large Higgs VEV $\langle H \rangle \approx 175$ GeV can lead to small neutrino masses providing that the mass scale M is high enough. E.g. if M is equal to the GUT scale $1.75 \cdot 10^{16}$ GeV then $m_{LL}^\nu = \lambda_\nu 1.75 \cdot 10^{-3}$ eV. To obtain larger neutrino masses we need to reduce M below the GUT scale (since we cannot make λ_ν too large otherwise it becomes non-perturbative).

Typically in physics whenever we see a large mass scale M associated with a non-renormalizable operator we tend to associate it with tree level exchange of some heavy particle or particles of mass M in order to make the high energy theory renormalizable once again. This idea leads directly to the see-saw mechanism where the exchanged particles can either couple to HL , in which case they must be either fermionic singlets (right-handed neutrinos) or fermionic triplets, or they can couple to LL and HH , in which case they must be scalar triplets. These three possibilities have been called the type I, III and II see-saw mechanisms, respectively. If the coupling λ_ν is very small (for some reason) then M could even be lowered to the TeV scale and the see-saw scale could be probed at the LHC ⁵⁾, however the see-saw mechanism then no longer solves the problem of the smallness of neutrino masses.

There are other ways to generate Majorana neutrino masses which lie outside of the above discussion. One possibility is to introduce additional Higgs singlets

and triplets in such a way as to allow neutrino Majorana masses to be generated at either one ⁶⁾ or two ⁷⁾ loops. Another possibility is within the framework of R-parity violating Supersymmetry in which the sneutrinos $\tilde{\nu}$ get small VEVs inducing a mixing between neutrinos and neutralinos χ leading to Majorana neutrino masses $m_{LL} \approx \langle \tilde{\nu} \rangle^2 / M_\chi$, where for example $\langle \tilde{\nu} \rangle \approx \text{MeV}$, $M_\chi \approx \text{TeV}$ leads to $m_{LL} \approx \text{eV}$. A viable spectrum of neutrino masses and mixings can be achieved at the one loop level ⁸⁾.

8. Normal or Inverted?

If the mass ordering is inverted then this may indicate a new symmetry such as $L_e - L_\mu - L_\tau$ ⁹⁾ or a $U(1)$ family symmetry ¹⁰⁾. However let us assume that the hierarchy is normal and proceed down the road map to the next experimental question.

9. Very precise tri-bimaximal mixing?

It is a striking fact that current data on lepton mixing is (approximately) consistent with the so-called tri-bimaximal (TB) mixing pattern ¹¹⁾,

$$U_{TB} = \begin{pmatrix} \sqrt{\frac{2}{3}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{6}} & -\frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \end{pmatrix} P_{Maj}, \quad (1)$$

where P_{Maj} is the diagonal phase matrix involving the two observable Majorana phases. However there is no convincing reason to expect exact TB mixing, and in general we expect deviations. These deviations can be parametrized by three parameters r, s, a defined as ¹²⁾:

$$\sin \theta_{13} = \frac{r}{\sqrt{2}}, \quad \sin \theta_{12} = \frac{1}{\sqrt{3}}(1 + s), \quad \sin \theta_{23} = \frac{1}{\sqrt{2}}(1 + a). \quad (2)$$

Global fits of the conventional mixing angles ^{13,14)} can be translated into the 1σ ranges

$$0.14 < r < 0.24, \quad -0.05 < s < 0.02, \quad -0.04 < a < 0.10. \quad (3)$$

Note in particular that the central value of r is now 0.2 which corresponds to a 2σ indication for a non-zero reactor angle as discussed at this meeting by Fogli ¹⁴⁾.

Clearly a non-zero value of r , if confirmed, would rule out TB mixing. However it is possible to preserve the good predictions that $s = a = 0$, by postulating a modified form of mixing matrix called tri-bimaximal-reactor (TBR) mixing ¹⁵⁾,

$$U_{TBR} = \begin{pmatrix} \sqrt{\frac{2}{3}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}}re^{-i\delta} \\ -\frac{1}{\sqrt{6}}(1 + re^{i\delta}) & \frac{1}{\sqrt{3}}(1 - \frac{1}{2}re^{i\delta}) & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{6}}(1 - re^{i\delta}) & -\frac{1}{\sqrt{3}}(1 + \frac{1}{2}re^{i\delta}) & \frac{1}{\sqrt{2}} \end{pmatrix} P_{Maj}. \quad (4)$$

Note that TBR mixing is distinct from the tri-maximal proposal ¹⁶⁾ that the second column of the mixing matrix should consist of a column with all elements equal to $1/\sqrt{3}$. On the one hand the TBR mixing proposal, to leading order in r , predicts the deviation parameters $s = a = 0$ for all r , whereas on the other hand the tri-maximal mixing proposal, to the same approximation, predicts $s = 0$ but $a = -(r/2)\cos\delta$. Thus, tri-bimaximal-reactor mixing and tri-maximal mixing may be distinguished by accurate determinations of a, r, δ (i.e. θ_{23}, θ_{13} and $\cos\delta$) at future high precision neutrino facilities.

10. Family Symmetry?

Assuming that TB or TBR mixing is very precise and is not an accident, it could be interpreted as a signal of an underlying family symmetry. Indeed I am unaware of any viable alternative at present. To understand the emergence of a family symmetry, let us expand the neutrino mass matrix in the diagonal charged lepton basis, assuming exact TB mixing, as $m_{LL}^\nu = U_{TB}\text{diag}(m_1, m_2, m_3)U_{TB}^T$ leading to (absorbing the Majorana phases in m_i):

$$m_{LL}^\nu = \frac{m_3}{2}\Phi_3\Phi_3^T + \frac{m_2}{3}\Phi_2\Phi_2^T + \frac{m_1}{6}\Phi_1\Phi_1^T \quad (5)$$

where $\Phi_3^T = (0, 1, 1)$, $\Phi_2^T = (1, 1, -1)$, $\Phi_1^T = (2, -1, 1)$ and m_i are the physical neutrino masses. This shows that the neutrino mass matrix corresponding to TB mixing may be constructed from the very simple orthogonal column vectors Φ_i , whose simplicity motivates an underlying non-Abelian family symmetry involving all three families. The idea is that Φ_i are promoted to new Higgs fields called “flavons” whose VEVs break the family symmetry, with the particular vacuum alignments as above. Such vacuum alignments can more readily be achieved if the non-Abelian family symmetry is a discrete symmetry containing a permutation symmetry capable of leading to $\langle |\Phi_2^T| \rangle \propto (1, 1, 1)$ ¹⁷⁾. A minimal choice of such family symmetry seems to be A_4 ¹⁸⁾ which only involves the flavon $\langle |\Phi_2^T| \rangle \propto (1, 1, 1)$ together with a further flavon $\langle |\Phi_0^T| \rangle \propto (0, 0, 1)$. Such minimal A_4 models lead to neutrino mass sum rules between the three masses m_i , resulting in/from a simplified mass matrix in Eq.5. A_4 may result from 6D orbifold models ¹⁹⁾.

It is possible to derive the TB form of the neutrino mass matrix in Eq.5 from the see-saw mechanism in a very elegant way as follows. In the diagonal right-handed neutrino mass basis we may write $M_{RR}^\nu = \text{diag}(M_A, M_B, M_C)$ and the Dirac mass matrix as $m_{LR}^\nu = (A, B, C)$ where A, B, C are three column vectors. Then the type I see-saw formula $m_{LL}^\nu = m_{LR}^\nu (M_{RR}^\nu)^{-1} (m_{LR}^\nu)^T$ gives

$$m_{LL}^\nu = \frac{AA^T}{M_A} + \frac{BB^T}{M_B} + \frac{CC^T}{M_C}. \quad (6)$$

By comparing Eq.6 to the TB form in Eq.5 it is clear that TB mixing will be achieved if $A \propto \Phi_3$, $B \propto \Phi_2$, $C \propto \Phi_1$, with each of $m_{3,2,1}$ originating from a particular

right-handed neutrino of mass $M_{A,B,C}$, respectively ²¹⁾. This mechanism allows a completely general neutrino mass spectrum and, since the resulting m_{LL}' is form diagonalizable, it is referred to as form dominance (FD) ²⁰⁾. For example, it has recently been show that the A_4 see-saw models ¹⁷⁾ satisfy FD ²⁰⁾.

If $m_1 \ll m_2 < m_3$ then the precise form of C becomes irrelevant, and in this case FD reduces to constrained sequential dominance (CSD)²¹⁾. The CSD mechanism has been applied in this case to models based on the family symmetries $SO(3)$ ^{21,22)} and $SU(3)$ ²³⁾, and their discrete subgroups ²⁴⁾.

It is possible to achieve TBR mixing, corresponding to $s = a = 0$ but $r \neq 0$, by a slight modification to the CSD conditions,

$$B = \frac{b}{\sqrt{3}} \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}, \quad A = \frac{c}{\sqrt{2}} \begin{pmatrix} \varepsilon \\ 1 \\ 1 \end{pmatrix}. \quad (7)$$

We refer to this as Partially Constrained Sequential Dominance (PCSD)¹⁵⁾, since one of the conditions of CSD is maintained, while the other one is violated by the parameter ε . Note that the introduction of the parameter ε also implies a violation of FD since the columns of the Dirac mass matrix A, B can no longer be identified with the columns of the MNS matrix, due to the non-orthogonality of A and B . To leading order in $|m_2|/|m_3|$ the mass matrix resulting from Eq.7 leads to TBR mixing where we identify ¹⁵⁾,

$$m_1 = 0, \quad m_2 = b^2/M_B, \quad m_3 = a^2/M_A, \quad \varepsilon = re^{-i\delta}. \quad (8)$$

Thus, the TBR form of mixing matrix in Eq.4 will result, to leading order in $|m_2|/|m_3|$.

11. Hierarchical or Degenerate?

This key experimental question may be decided by the same experiments as will also determine the nature of neutrino mass (Dirac or Majorana) ^{?)}. Although not a theorem, it seems that a hierarchical spectrum could indicate a type I see-saw mechanism, while a (quasi) degenerate spectrum could imply a type II see-saw mechanism. It is possible that a type II see-saw mechanism could naturally explain the degenerate mass scale with the degeneracy enforced by an $SO(3)$ family symmetry, while the type I see-saw part could be responsible for the small neutrino mass splittings and the (TB) mixing ²⁵⁾. An A_4 model of quasi-degenerate neutrinos with TB mixing, working at the effective neutrino mass operator level, was considered recently in ²⁶⁾.

12. GUTs and/or Strings?

Finally we have reached the end of the decision tree, with the possibility of an all-encompassing unified theory of flavour based on GUTs and/or strings. Such theories

could also include a family symmetry in order to account for the TB mixing. There are many possibilities for the choice of family symmetry and GUT symmetry. Examples include the Pati-Salam gauge group $SU(4)_{PS} \times SU(2)_L \times SU(2)_R$ in combination with $SU(3)$ ²³⁾, $SO(3)$ ^{21,22)}, A_4 ²⁷⁾ or Δ_{27} ²⁸⁾. Other examples are based on $SU(5)$ GUTs in combination with A_4 ²⁹⁾ or T' ³⁰⁾.

For example, it is straightforward to implement the above example of PCSD into realistic GUT models with non-Abelian family symmetry spontaneously broken by flavons which are based on the CSD mechanism ^{21,23,22)}. In such models the columns of the Dirac mass matrix in the diagonal right-handed neutrino mass basis are determined by flavon vacuum alignment, with the column B identified with a triplet flavon ϕ_{123} and the column A identified with a triplet flavon ϕ_{23} and it is quite easy to obtain a correction to the vacuum alignment such that

$$\langle \phi_{123} \rangle = \frac{b}{\sqrt{3}} \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}, \quad \langle \phi_{23} \rangle = \frac{a}{\sqrt{2}} \begin{pmatrix} \varepsilon \\ 1 \\ 1 \end{pmatrix}, \quad (9)$$

in direct correspondence with Eq.7. For example, in such models based on the discrete family symmetry A_4 ²²⁾, the flavon vacuum expectation value (VEV) $\langle \phi_{123} \rangle$ will preserve a Z_2 subgroup of the original discrete family symmetry corresponding to an A_4 generator S ¹⁹⁾, while the flavon VEV $\langle \phi_{23} \rangle$ will violate this subgroup even in the limit that $\varepsilon = 0$. It is therefore natural to assume some misalignment of $\langle \phi_{23} \rangle$ since, unlike $\langle \phi_{123} \rangle$, it is not protected by any symmetry.

In typical Family Symmetry $\otimes$ GUT models the origin of the quark mixing angles derives predominantly from the down quark sector, which in turn is closely related to the charged lepton sector. In order to reconcile the down quark and charged lepton masses, simple ansatze, such as the Georgi-Jarlskog hypothesis ³¹⁾, lead to very simple approximate expectations for the charged lepton mixing angles such as $\theta_{12}^e \approx \lambda/3$, $\theta_{23}^e \approx \lambda^2$, $\theta_{13}^e \approx \lambda^3$, where $\lambda \approx 0.22$ is the Wolfenstein parameter from the quark mixing matrix. If the family symmetry enforces accurate TB mixing in the neutrino sector, then $\theta_{12}^e \approx \lambda/3$ charged lepton corrections will cause deviations from TB mixing in the physical lepton mixing angles, and lead to a sum rule relation ^{21,32,33)}, which can be conveniently expressed as ¹²⁾ $s \approx r \cos \delta$ where $r \approx \lambda/3$ and δ is the observable CP violating oscillation phase, with RG corrections of less than one degree ³⁴⁾. Such sum rules can be tested in future high precision neutrino oscillation experiments ³⁵⁾.

Note that in such a GUT-flavour framework, one expects the charged lepton corrections to the neutrino mixing angles to be less than of order $\theta_{12}^e/\sqrt{2}$ (where typically θ_{12}^e is a third of the Cabibbo angle) plus perhaps a further 1° from renormalization group (RG) corrections. Thus such theoretical corrections cannot account for an observed reactor angle as large as 8° , corresponding to $r = 0.2$, starting from the

hypothesis of exact TB neutrino mixing.

13. Conclusion

Neutrino mass and mixing clearly requires new physics beyond the SM, but in which direction should we go? There are many roads for model building, but we have seen that answers to key experimental questions will provide the sign posts *en route* to a unified theory of flavour.

In particular we would like to emphasize that a measurement of a large reactor angle, consistent with the present 2σ indication for $r = 0.2$, can still be consistent with tri-bimaximal solar and atmospheric mixing, corresponding to $s = a = 0$, according to the tri-bimaximal-reactor mixing hypothesis. By contrast, tri-maximal mixing predicts $s = 0$ but $a = -(r/2) \cos \delta$.

Alternatively the presence of a large reactor angle could be a sign that TB mixing is the wrong starting point and instead it is better to start from bi-maximal neutrino mixing and then invoke large charged lepton corrections to correct the reactor and solar angles, as discussed by Altarelli at this meeting³⁶).

The common feature of all these approaches is the presence of an underlying family symmetry, even though the reactor angle may be quite large. The existence of such disparate approaches only underlines the need for further high precision data from the neutrino experiments in order to resolve which approach is correct.

14. Acknowledgements

I would like to thank the organizers of the XIII International Workshop on Neutrino Telescopes, Venice, March 10-13, 2009, Venice for their kind hospitality, and for partial support from the following grants: STFC Rolling Grant ST/G000557/1; EU Network MRTN-CT-2004-503369; EU ILIAS RII3-CT-2004-506222.

15. References

- 1) N. Arkani-Hamed, S. Dimopoulos, G. R. Dvali and J. March-Russell, Phys. Rev. D **65** (2002) 024032 [arXiv:hep-ph/9811448].
- 2) S. Antusch, O. J. Eyton-Williams and S. F. King, JHEP **0508** (2005) 103 [arXiv:hep-ph/0505140].
- 3) For an overview with references see e.g. S. J. Huber, arXiv:hep-ph/0211056.
- 4) P. Fileviez Perez, T. Han, G. y. Huang, T. Li and K. Wang, Phys. Rev. D **78** (2008) 015018 [arXiv:0805.3536 [hep-ph]].
- 5) S. F. King and T. Yanagida, Prog. Theor. Phys. **114** (2006) 1035 [arXiv:hep-ph/0411030].
- 6) A. Zee, Phys. Lett. B **93** (1980) 389 [Erratum-ibid. B **95** (1980) 461].

- 7) K. S. Babu, Phys. Lett. B **203** (1988) 132.
- 8) See for example: M. Hirsch, M. A. Diaz, W. Porod, J. C. Romao and J. W. F. Valle, Phys. Rev. D **62** (2000) 113008 [Erratum-ibid. D **65** (2002) 119901] [arXiv:hep-ph/0004115].
- 9) S. T. Petcov, Phys. Lett. B **110** (1982) 245.
- 10) S. F. King and N. N. Singh, Nucl. Phys. B **596** (2001) 81 [arXiv:hep-ph/0007243].
- 11) P. F. Harrison, D. H. Perkins and W. G. Scott, Phys. Lett. B **530** (2002) 167 [arXiv:hep-ph/0202074].
- 12) S. F. King, Phys. Lett. B **659** (2008) 244 [arXiv:0710.0530 [hep-ph]].
- 13) M. Maltoni, talk at XIII International Workshop on Neutrino Telescopes, Venice, March 10-13, 2009, Venice; M. Maltoni, T. Schwetz, M. A. Tortola and J. W. F. Valle, New J. Phys. **6** (2004) 122 [arXiv:hep-ph/0405172]; T. Schwetz, M. Tortola and J. W. F. Valle, arXiv:0808.2016 [hep-ph].
- 14) G. Fogli, talk at XIII International Workshop on Neutrino Telescopes, Venice, March 10-13, 2009, Venice; G. L. Fogli, E. Lisi, A. Marrone, A. Palazzo and A. M. Rotunno, Phys. Rev. Lett. **101** (2008) 141801 [arXiv:0806.2649 [hep-ph]].
- 15) S. F. King, arXiv:0903.3199 [hep-ph].
- 16) W. G. Scott, talk at XIII International Workshop on Neutrino Telescopes, Venice, March 10-13, 2009, Venice; L. Wolfenstein, Phys. Rev. D **18** (1978) 958; P. F. Harrison and W. G. Scott, Phys. Lett. B **535** (2002) 163 [arXiv:hep-ph/0203209]; J. D. Bjorken, P. F. Harrison and W. G. Scott, Phys. Rev. D **74** (2006) 073012 [arXiv:hep-ph/0511201]; X. G. He and A. Zee, Phys. Lett. B **645** (2007) 427 [arXiv:hep-ph/0607163]; W. Grimus and L. Lavoura, JHEP **0809** (2008) 106 [arXiv:0809.0226 [hep-ph]]; C. H. Albright and W. Rodejohann, arXiv:0812.0436 [hep-ph].
- 17) G. Altarelli and F. Feruglio, Nucl. Phys. B **720** (2005) 64 [arXiv:hep-ph/0504165].
- 18) E. Ma and G. Rajasekaran, Phys. Rev. D **64** (2001) 113012 [arXiv:hep-ph/0106291].
- 19) G. Altarelli, F. Feruglio and Y. Lin, Nucl. Phys. B **775** (2007) 31 [arXiv:hep-ph/0610165].
- 20) M. C. Chen and S. F. King, arXiv:0903.0125 [hep-ph].
- 21) S. F. King, JHEP **0508** (2005) 105 [arXiv:hep-ph/0506297].
- 22) S. F. King and M. Malinsky, JHEP **0611** (2006) 071 [arXiv:hep-ph/0608021].
- 23) I. de Medeiros Varzielas and G. G. Ross, Nucl. Phys. B **733** (2006) 31 [arXiv:hep-ph/0507176].
- 24) I. de Medeiros Varzielas, S. F. King and G. G. Ross, Phys. Lett. B **644** (2007) 153 [arXiv:hep-ph/0512313].
- 25) S. Antusch and S. F. King, Nucl. Phys. B **705** (2005) 239

- [arXiv:hep-ph/0402121].
- 26) I. d. M. Varzielas, G. G. Ross and M. Serna, arXiv:0811.2226 [hep-ph].
 - 27) S. F. King and M. Malinsky, Phys. Lett. B **645** (2007) 351 [arXiv:hep-ph/0610250].
 - 28) I. de Medeiros Varzielas, S. F. King and G. G. Ross, Phys. Lett. B **648** (2007) 201 [arXiv:hep-ph/0607045].
 - 29) G. Altarelli, F. Feruglio and C. Hagedorn, JHEP **0803** (2008) 052 [arXiv:0802.0090 [hep-ph]].
 - 30) M. C. Chen and K. T. Mahanthappa, Phys. Lett. B **652** (2007) 34 [arXiv:0705.0714 [hep-ph]].
 - 31) H. Georgi and C. Jarlskog, Phys. Lett. B **86** (1979) 297.
 - 32) I. Masina, Phys. Lett. B **633** (2006) 134 [arXiv:hep-ph/0508031];
 - 33) S. Antusch and S. F. King, Phys. Lett. B **631** (2005) 42 [arXiv:hep-ph/0508044];
 - 34) S. Boudjemaa and S. F. King, arXiv:0808.2782 [hep-ph].
 - 35) S. Antusch, P. Huber, S. F. King and T. Schwetz, JHEP **0704** (2007) 060 [arXiv:hep-ph/0702286].
 - 36) G. Altarelli, talk at XIII International Workshop on Neutrino Telescopes, Venice, March 10-13, 2009, Venice; G. Altarelli, F. Feruglio and L. Merlo, arXiv:0903.1940 [hep-ph]; S. Antusch, S. F. King and R. N. Mohapatra, Phys. Lett. B **618** (2005) 150 [arXiv:hep-ph/0504007].